Rehabilitation Protocol: Arthroscopic Posterior Shoulder Stabilization



Signature:

Craig M. Capeci, MD

Orthopaedic Surgery & Sports Medicine Clinical Assistant Professor __1095 Park Ave, NY 212-427-7750 __1056 5th Ave, NY 212-348-3636 __263 7th Ave, Bklyn 718-246-8700

Date: _____

| Name: | Date: |
|--|---|
| Diagnosis: | Date of Surgery: |
| rehab under guidance of PT Range of Motion – None for Weeks 0-3 Weeks 3-6: Begin passive ROM - Restrict n 45° of Internal Rotation Therapeutic Exercise Elbow/Wrist/Hand Range of Motion Grip Strengthening | ction and 0° of rotation) except for showering and notion to 90° of Forward Flexion, 90° of Abduction, and ities: Codman's, Anterior Capsule Mobilization |
| | otator Cuff Isometrics s for Rotator Cuff/Scapular Stabilizers/Biceps and elow the horizontal plane during this phase – utilize |
| Phase III (Weeks 12-16) Range of Motion – Progress to full AROM without of the Therapeutic Exercise – Advance Phase II exercises Emphasize Glenohumeral Stabilization, Exton Begin UE ergometer/endurance activities Modalities per PT discretion | |
| Phase IV (Months 4-6) Range of Motion – Full without discomfort Therapeutic Exercise – Continue with strengthening Sport/Work specific rehabilitation – Plyom Continue with endurance activities Return to sports at 6 months if approved Modalities per PT discretion | |
| Comments: | |
| Frequency: times per week Dur | ation: weeks |