Rehabilitation Protocol: Arthroscopic Anterior Shoulder Stabilization (Bankart Repair)



Craig M. Capeci, MD

Orthopaedic Surgery & Sports Medicine Clinical Assistant Professor __1095 Park Ave, NY 212-427-7750 __1056 5th Ave, NY 212-348-3636 __263 7th Ave, Bklyn 718-246-8700

Name:	Date:
Diagnosis:	Date of Surgery:
 Phase I (Weeks 0-4) Sling immobilization at all times except for showering and rehab under guidance of PT Range of Motion – PROM → AAROM → AROM as tolerated Restrict motion to 90° of Forward Flexion, 45° of Abduction, 30° of External Rotation and Internal Rotation to stomach Therapeutic Exercise Elbow/Wrist/Hand Range of Motion Grip Strengthening Isometric Abduction, Internal/External Rotation exercises with elbow at side Heat/Ice before and after PT sessions 	
 Phase II (Weeks 4-6) Discontinue sling immobilization Range of Motion – Increase Forward Flexion, Internated Therapeutic Exercise Advance isometrics from Phase I to use of a to Continue with Elbow/Wrist/Hand Range of Degin Prone Extensions and Scapular Stability Gentle joint mobilization Modalities per PT discretion 	heraband within AROM limitations
 Phase III (Weeks 6-12) Range of Motion – Progress to full AROM without discomfort Therapeutic Exercise – Advance theraband exercises to light weights (1-5 lbs) 8-12 repetitions/2-3 sets for Rotator Cuff, Deltoid and Scapular Stabilizers Continue and progress with Phase II exercises Begin UE ergometer Modalities per PT discretion 	
 Phase IV (Months 3-6) Range of Motion – Full without discomfort Therapeutic Exercise – Advance exercises in Phase III (strengthening 3x per week) Sport/Work specific rehabilitation Return to throwing at 4.5 months Return to sports at 6 months if approved Modalities per PT discretion 	
Comments:	
Frequency: times per week Dura	tion: weeks