Rehabilitation Protocol: Arthroscopic Meniscectomy/Chondral Debridement



Craig M. Capeci, MD

Orthopaedic Surgery & Sports Medicine Clinical Assistant Professor __1095 Park Ave, NY 212-427-7750 __1056 5th Ave, NY 212-348-3636 __263 7th Ave, Bklyn 718-246-8700

Name:	Date:
Diagnosis:	Date of Surgery:
 Phase I (Weeks 0-2) Weightbearing: As tolerated with crutches (for a sample of Motion – AAROM → AROM as tolerated of Goal: Immediate full range of motion Therapeutic Exercises Quad and Hamstring sets Heel slides Co-contractions Isometric adduction and abduction exercitions Straight-leg raises Patellar mobilization 	i
 Phase II (Weeks 2-4) Weightbearing: As tolerated Range of Motion – maintain full ROM – gentle parameters Therapeutic Exercises Quadriceps and Hamstring strengthening Lunges Wall-sits Balance exercises – Core work 	
Phase III (Weeks 4-6) • Weightbearing: Full weightbearing • Range of Motion – Full/Painless ROM • Therapeutic Exercises • Leg press • Hamstring curls • Squats • Plyometric exercises • Endurance work • Return to athletic activity as tolerated	
Comments:	
Frequency: times per week Du	ration: weeks
Signature	Date